arXiv:1612.05104v1 [math.PR] 8 Dec 2016

AN APPROACH THEORETIC VERSION OF ANSCOMBE'S THEOREM

BEN BERCKMOES

Abstract. We establish an approach theoretic version of Anscombe's theorem, which enhances its applicational power.

1. Introduction

In Anscombe theory the stability of weak convergence of random variables under index randomization is investigated. Let us explain this idea more precisely.

Let X be a separable metric space with metric d, and consider X-valued random variables ξ and $(\xi_n)_n$. We let $\stackrel{w}{\to}$ stand for weak convergence, i.e. $\xi_n \stackrel{w}{\to} \xi$ means that, for all $f : X \to \mathbb{R}$ bounded and continuous, $\mathbb{E}[f(\xi_n)] \to \mathbb{E}[f(\xi)]$. Furthermore, consider a sequence $(N_n)_n$ of $\mathbb{N}$-valued random variables. Anscombe theory deals with the following natural question. Under which conditions does the implication $\xi_n \stackrel{w}{\to} \xi \Rightarrow \xi_{N_n} \stackrel{w}{\to} \xi$ hold?

Let $\stackrel{\mathbb{P}}{\to}$ stand for convergence in probability, i.e. $\xi_n \stackrel{\mathbb{P}}{\to} \xi$ means that, for each $\epsilon > 0$, $\mathbb{P}[d(\xi, \xi_n) \geq \epsilon] \to 0$. Furthermore, we say that a sequence $(\xi_n)_n$ of random variables satisfies Anscombe's condition iff for each $\epsilon > 0$ there exist $\delta > 0$ and n_0 such that $\mathbb{P}\left[\max_{m=\lceil (1-\delta)n \rceil}^{\lfloor (1+\delta)n \rfloor} d(\xi_n, \xi_m) \geq \epsilon\right] < \epsilon$ for all $n \geq n_0$, $\lfloor \cdot \rfloor$ being the floor function, and $\lceil \cdot \rceil$ the ceiling function. This condition guarantees that a sequence of random variables 'oscillates slowly'.

The importance of Anscombe's condition is reflected by the following result, originally obtained by Anscombe in [A52], which roughly states that if the random sequence $(N_n)_n$ is eventually close to a deterministic sequence $(k_n)_n$ which tends to ∞, and $(\xi_n)_n$ satisfies Anscombe's condition, then weak convergence of $(\xi_n)_n$ to ξ implies weak convergence of $(\xi_{N_n})_n$ to ξ.

Theorem 1.1 (Anscombe)**.** *Let ξ and $(\xi_n)_n$ be X-valued random variables and $(N_n)_n$ $\mathbb{N}$-valued random variables. Suppose, in addition, that*

(a) *there exists $(k_n)_n$ in $\mathbb{R}_0^+$ such that $k_n \to \infty$ and $\frac{N_n}{k_n} \stackrel{\mathbb{P}}{\to} 1$,*
(b) *$(\xi_n)_n$ satisfies Anscombe's condition.*

Then $\xi_n \stackrel{w}{\to} \xi \Rightarrow \xi_{N_n} \stackrel{w}{\to} \xi$.

A beautiful review of the rich history of Ancombe theory is given in [G12]. We wish to point out that condition (a) in Anscombe's theorem actually 'kills' the randomness of $(N_n)_n$, which makes the result often inapplicable

Key words and phrases. Anscombe's theorem, approach theory.
Ben Berckmoes is post doctoral fellow at the Fund for Scientific Research of Flanders (FWO).

(see e.g. [BL13], subsection 7.1.1, paragraph 2, and [BM16]). We will establish an approach theoretic version of Anscombe's theorem, which enhances its applicational power, in the next section.

2. An approach theoretic version of Anscombe's theorem

Define, for any sequence $(\xi_n)_n$ of X-valued random variables, the Anscombe index as

$$\chi_{\text{Ansc}}\left\langle (\xi_n)_n \right\rangle = \sup_{\epsilon>0} \inf_{\delta>0} \limsup_{n\to\infty} \mathbb{P}\left[\max_{m=\lceil (1-\delta)n \rceil}^{\lfloor (1+\delta)n \rfloor} d(\xi_n, \xi_m) \geq \epsilon \right].$$

Then $\chi_{\text{Ansc}}\left\langle (\xi_n)_n \right\rangle$ takes values between 0 and 1 and it is 0 if and only if $(\xi_n)_n$ satisfies Anscombe's condition.

Furthermore, recall that λ_w and $\lambda_{\mathbb{P}}$ are the limit operators for respectively the weak approach structure and the approach structure of convergence in probability ([L15]). That is, for X-valued random variables ξ and $(\xi_n)_n$,

$$\begin{aligned}
&\lambda_w(\xi_n \to \xi) \\
&\quad = \sup_{f\in\mathcal{C}(X,[0,1])} \limsup_{n\to\infty} \left|\mathbb{E}\left[f(\xi) - f(\xi_n)\right]\right| \\
&\quad = \sup_{\alpha\in\mathbb{R}_0^+} \limsup_{n\to\infty} \sup_{A \text{ Borel}} \left(\mathbb{P}[\xi \in A] - \mathbb{P}\left[\xi_n \in A^{(\alpha)}\right]\right) \\
&\quad = \sup_{G \text{ open}} \limsup_{n\to\infty} \left(\mathbb{P}[\xi \in G] - \mathbb{P}[\xi_n \in G]\right) \\
&\quad = \sup_{F \text{ closed}} \limsup_{n\to\infty} \left(\mathbb{P}[\xi_n \in F] - \mathbb{P}[\xi \in F]\right) \\
&\quad = \sup_{A\in\mathcal{A}_\xi} \limsup_{n\to\infty} \left|\mathbb{P}[\xi \in A] - \mathbb{P}[\xi_n \in A]\right|,
\end{aligned}$$

with $\mathcal{C}(X,[0,1])$ the set of continuous maps of X into $[0,1]$ and $\mathcal{A}_\xi$ the collection of Borel sets $A \subset X$ for which the $\mathbb{P}$-probability that ξ is on the boundary of A is 0, and

$$\lambda_{\mathbb{P}}(\xi_n \to \xi) = \sup_{\epsilon>0} \limsup_{n\to\infty} \mathbb{P}\left[d(\xi, \xi_n) \geq \epsilon\right].$$

Notice that $\lambda_w(\xi_n \to \xi)$ (respectively $\lambda_{\mathbb{P}}(\xi_n \to \xi)$) is a number between 0 and 1 which measures how far $(\xi_n)_n$ deviates from converging weakly (respectively in probability) to ξ.

In this framework, the following result holds. A proof is given in the next section.

Theorem 2.1 (Approach theoretic Anscombe)**.** *Let ξ and $(\xi_n)_n$ be X-valued random variables and $(N_n)_n$ $\mathbb{N}$-valued random variables. Then*

$$\lambda_w\left(\xi_{N_n} \to \xi\right) \leq \lambda_w\left(\xi_n \to \xi\right) + \chi_{\text{Ansc}}\left\langle (\xi_n)_n \right\rangle + \inf_{(k_n)_n} \lambda_{\mathbb{P}}\left(\frac{N_n}{k_n} \to 1\right), \quad (1)$$

the infimum being taken over all $(k_n)_n$ in $\mathbb{R}_0^+$ with $k_n \to \infty$.

Notice that Theorem 2.1 is a strict generalization of Anscombe's theorem (Theorem 1.1). Indeed, if the conditions in Anscombe's theorem are fulfilled, then the right-hand side in (1) becomes 0, whence $\lambda_w\left(\xi_{N_n} \to \xi\right) = 0$, and we conclude that $\xi_{N_n} \xrightarrow{w} \xi$. Theorem 2.1 has the advantage that even in cases

where the conditions in Anscombe's theorem fail to be fulfilled, it continues to provide valuable results. Applications will be given in subsequent work.

3. Proof of Theorem 2.1

Let, for positive numbers ζ, η and θ, and $(k_n)_n$ in $\mathbb{R}_0^+$ with $k_n \to \infty$, $\lambda_w(\xi_n \to \xi) < \zeta$, $\chi_{\text{Ansc}}\langle(\xi_n)_n\rangle < \eta$, and $\lambda_{\mathbb{P}}\left(\frac{N_n}{k_n} \to 1\right) < \theta$. Fix $\epsilon > 0$ and $F \subset X$ closed. Then there exist $\delta > 0$ and n_1 such that for all $n \geq n_1$

$$\mathbb{P}\left[\max_{m=\lceil(1-\delta)n\rceil}^{\lfloor(1+\delta)n\rfloor} d(\xi_n, \xi_m) \geq \epsilon\right] < \eta.$$

Also, there exists n_2 such that $k_n \geq n_1$ for all $n \geq n_2$. Furthermore, there exists n_3 such that for all $n \geq n_3$

$$\mathbb{P}[|k_n - N_n| > \delta k_n] < \theta.$$

Put $n_0 = \max\{n_1, n_2, n_3\}$. Now assume without loss of generality that $(k_n)_n$ is integer-valued. Then, for $n \geq n_0$,

$$\begin{aligned}
&\mathbb{P}\left[\xi_{N_n} \in F\right] - (\eta + \theta) \\
&\quad\leq \mathbb{P}\left[\{\xi_{N_n} \in F\} \cap \left\{\max_{m=\lceil(1-\delta)k_n\rceil}^{\lfloor(1+\delta)k_n\rfloor} d(\xi_{k_n}, \xi_m) < \epsilon\right\} \cap \{|k_n - N_n| \leq \delta k_n\}\right] \\
&\quad\leq \mathbb{P}\left[\xi_{k_n} \in F^{(\epsilon)}\right],
\end{aligned}$$

whence

$$\limsup_{n\to\infty} \mathbb{P}\left[\xi_{N_n} \in F\right] \leq \limsup_{n\to\infty} \mathbb{P}\left[\xi_{k_n} \in F^{(\epsilon)}\right] + \eta + \theta \leq \mathbb{P}\left[\xi \in F^{(\epsilon)}\right] + \zeta + \eta + \theta.$$

This finishes the proof of (1).

References

[A52] Anscombe, F. J. *Large-sample theory of sequential estimation.* Proc. Cambridge Philos. Soc. 48, (1952). 600–607.

[BL13] Bartroff, J.; Lai, T. L.; Shih, M.-C. *Sequential experimentation in clinical trials. Design and analysis.* Springer Series in Statistics. Springer, New York, 2013.

[BM16] Berckmoes, B.; Molenberghs, G. *On the asymptotic normality and the construction of confidence intervals for estimators after sampling with probabilistic and deterministic stopping rules* (2016), https://arxiv.org/abs/1608.02801.

[L15] Lowen, R. *Index analysis. Approach theory at work.* Springer Monographs in Mathematics. Springer, London, 2015.

[G12] Gut, A. *Anscombe's theorem 60 years later.* Sequential Anal. 31 (2012), no. 3, 368–396.